



**Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking**

Fayet-Moore, Flavia; McConnell, Andrew; Petocz, Peter; Cassettari, Tim; Tuck, Kate; Blumfield, Michelle ; Warner, Molly; Marshall, Skye

*Published in:*  
Nutrition and Dietetics

*DOI:*  
[10.1111/1747-0080.12648](https://doi.org/10.1111/1747-0080.12648)

*Licence:*  
Other

[Link to output in Bond University research repository.](#)

*Recommended citation(APA):*  
Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2021). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, 78(2), 154-164. <https://doi.org/10.1111/1747-0080.12648>

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

# **The contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: a comparison using an objective versus subjective definition for snacking**

Running heading: Dietary snacking patterns of Australian adults

Flavia Fayet-Moore<sup>1\*</sup>, Andrew McConnell<sup>2</sup>, Peter Petocz<sup>3</sup>, Tim Cassettari<sup>4</sup>, Kate Tuck<sup>5</sup>,

Michelle Blumfield<sup>6</sup>, Molly Warner<sup>7</sup>, Skye Marshall<sup>8</sup>

1. PhD, APD, Nutrition Research Australia, Sydney, New South Wales, Australia.
2. BSc (AdvMaths), Nutrition Research Australia, Sydney, New South Wales, Australia.
3. PhD, Nutrition Research Australia, Sydney, New South Wales, Australia.
4. Bsc(Hons), APD, Nutrition Research Australia, Sydney, New South Wales, Australia.
5. BAppSci(Biophysics), Nutrition Research Australia, Sydney, New South Wales, Australia.
6. BNutrDiet (Hons), PhD, APD. Nutrition Research Australia, New South Wales, Australia.
7. MNutrDietPrac, APD, Nutrition Research Australia, Sydney, New South Wales, Australia.
8. PhD, APD, Nutrition Research Australia, Sydney, New South Wales, Australia; and Bond University Nutrition and Dietetics Research Group, Faculty of Health Sciences and Medicine, Bond University, Gold Coast, Queensland, Australia.

\*Corresponding Author: Skye Marshall, [skye@nraus.com](mailto:skye@nraus.com), Level 10, 20 Martin Place Sydney, New South Wales, Australia. Phone: +61 415 990 050.

**Manuscript words:** 3,929 (including references)

**Abstract words:** 247

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

## Abstract

**Aim:** There is limited information on the snacking behaviour of Australian adults, and the role of snacking in the diet may depend on how it is defined. This study aimed to compare the dietary snacking behaviours and associated nutritional intake and body composition in Australian adults, using an objective versus a subjective definition for snacking.

**Methods:** Cross-sectional data were analysed from the 2011-12 National Nutrition and Physical Activity Survey (n = 8,361, 19+ years). Objective snacking was defined based on time of day between main meals. Subjective snacking was self-reported by participants.

**Results:** Using the objective definition, 88.2% of adults were snack consumers; where snacking contributed 20.0% (SD: 20.0%) of total daily energy intake and 27.0% (SD 31.4%) of total daily discretionary energy. 41.3% (SD 37.1%) of snacking energy intake came from discretionary foods. Using the subjective definition, 98.5% of adults were snack consumers, where discretionary foods contributed 52.6% (SD 35.2%) of all snacking energy. The proportion of objective and subjective snacking energy from discretionary foods did not differ across BMI groups.

**Conclusions:** Objective snacking energy contributed more to core food groups than discretionary, was not associated with anthropometric measures, and contributed less to total discretionary energy than the evening meal. When snacking was defined subjectively, more than half of snacking energy was discretionary, suggesting that adults are more likely to perceive discretionary foods as snacks. Differences between snacking definitions means that associations between self-reported snacking and diet or health outcomes, should be interpreted with caution.

**Keywords:** nutrition assessment; energy intake; cross-sectional studies; diet, Western; snacks

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.  
This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

## Introduction

Globally, the dietary pattern of snacking is becoming increasingly popular. Snacking prevalence in adults ranges from 35% in China <sup>1</sup>, to 73% in Mexico, 74% in Brazil <sup>2,3</sup>, 77% in Canada <sup>4</sup> and up to 97% in the United States of America (USA) <sup>5</sup>. Although there is no consensus on the relationship between snacking and health <sup>6</sup>, snacking is often perceived as synonymous with consumption of highly processed or discretionary foods and beverages <sup>7,8</sup>, poor diet quality <sup>9</sup> and weight gain <sup>10-12</sup>. As such, there have been recommendations that snacking frequency be reduced <sup>7,8</sup>. However, snacking appears to facilitate energy balance in individuals with a recommended body mass index (BMI) (18.5 to 24.9 kg/m<sup>2</sup>) <sup>13</sup> and can add beneficial nutrients to the diet <sup>14</sup>.

There is inconsistency in how snack foods and snacking are defined. A snack food can be defined by its nutritional composition, time of consumption or be self-reported by the consumer <sup>15,16</sup>. Snacking behaviour may be defined by the time of consumption, the contribution to total energy intake, meal frequency, reported eating occasion, a combination of nutritional composition and temporal patterns, or whether it was motivated by hunger <sup>5,10,15</sup>. The lack of a universal definition impacts the interpretation of the snacking literature and limits development of evidence-based snacking dietary recommendations <sup>15,16</sup>.

There is limited information on the snacking behaviour of Australian adults, and the prevalence and contribution of snacking to the Australian adult diet is unknown. With nearly two thirds of Australian adults classified as overweight or obese according to

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

BMI<sup>17</sup> and a high intake of discretionary foods<sup>18,19</sup>, there is a need to investigate snacking in the Australian population and the associations that snacking has with the types of foods and beverages that are consumed, their contribution to energy and nutrient intakes, and the relationship with age, sex, and measures of body composition. Given the lack of a universal definition for snacking, there is also a need to determine the differences between definitions. Therefore, this study aimed to compare the dietary snacking behaviours and associated nutritional intake and body composition in Australian adults, using an objective versus subjective snacking definition.

## Methods

This cross-sectional study has been reported according to the STROBE statement<sup>20</sup>.

Cross-sectional data from the 2011-12 National Nutrition and Physical Activity Survey (NNPAS) were used. The NNPAS is a nationally representative survey that was part of the 2011-13 Australian Health Survey conducted by the Australian Bureau of Statistics (ABS)<sup>18</sup>, the methodology of which has been reported in detail elsewhere<sup>21</sup>. Briefly, 9341 adult (19-years or older) survey participants had their dietary intake assessed using an Automated Multiple-Pass Method to capture all foods and beverages consumed in the 24 hours prior to the interview day. Due to differences in meal patterns between weekdays and weekends, only weekdays were included<sup>22-24</sup>; therefore, a total of 8361 adults were included in the current analysis. The interview components of the survey were conducted under the Census and Statistics Act 1905 and therefore ethics approval was not necessary.

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

Objective and subjective definitions to measure snacking were used. For the objective definition, daily energy contribution was charted at half-hour intervals and meal periods began at the first energy increase and ended in the trough <sup>25</sup>. The three largest peaks were classified as main meals and the three smaller peaks were classified as between main meals (snacking time periods). Snacking time periods were thus classified as morning: 9.30am – 12.00pm, afternoon: 2.30pm – 5.30pm, and late night: 9.30pm – 5.30am (Figure 1). A snacking occasion was defined as the consumption of one or more food or beverage items if the consumption fell within the snacking time periods. For the subjective definition, participant defined snacking occasions were used. During the dietary recall interviews, respondents classified the reported eating occasion (REO) for each food and beverage consumed from a list of 11 pre-defined options which included four main meals (breakfast, brunch, lunch and dinner) and seven subjective snacking eating occasions (morning tea, afternoon tea, supper, snack, beverage/drink, extended consumption and other).

Dietary intake was calculated by the ABS using the 2011-13 Australian Food, Supplement and Nutrient Database (AUSNUT) developed by Food Standards Australia New Zealand (FSANZ) <sup>26</sup>. The contribution to total snacking energy from each sub-major food group, the proportion of consumers who reported snacking, and the portion size in grams per consumer of each sub-major food group was calculated. The mean daily energy intake and energy intake during each snacking time period (objective definition), the contribution from energy during snacking to total daily energy intake, the contribution from snacking to protein, fat, carbohydrate and total and added sugars were also calculated. Discretionary foods and beverages were

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

defined according to the Australian Dietary Guidelines which are foods and drinks not necessary to provide the nutrients the body needs, and are those high in saturated fats, added sugars, added salt, or alcohol and low in dietary fibre <sup>27</sup>. Discretionary foods in the survey were coded as discretionary by the ABS. The total daily discretionary energy intake, discretionary intake during snacking, contribution from snacking to total daily discretionary energy and contribution of discretionary energy to total snacking energy were calculated.

Respondents' anthropometric measures, including weight, height and waist circumference, were measured during NNPAS interview. BMI was calculated as kg/m<sup>2</sup>, and participants' BMI and waist circumference were categorised according to the World Health Organisation <sup>28</sup>.

The statistical package SPSS version 23.0 (IBM Corp., Armonk, NY, USA) was used for all analyses, with a p-value <0.001 treated as significant. General linear models were used to investigate the effects of a standard set of predictor variables (age group, sex, their interaction, total energy intake and BMI group); the response variables included proportion of energy from snacking, proportion of discretionary energy from snacking, proportion of energy from discretionary foods and number of snacking occasions. ANOVA tables and Chi-squared tests were used to indicate significant variables, and Bonferroni post-hoc analyses were performed to show pairwise significance between categories of respondents. Dependent t-tests were used to determine significant differences between the objective and subjective snacking definitions. Confidence intervals have been used for population proportions to show the precision of the result

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

and the sampling error. Standard deviations have been provided for population means to indicate the variance of the sample.

## Results

Of the 8361 adults, the majority were female (51.1%), aged 19-50 years (59.8%), and had a BMI  $>30\text{kg/m}^2$  (62.5%) (Table 1). The majority of adults were objective snack consumers (88.2%) and the highest prevalence of objective snacking was among those aged 31-50-years (89.8%;  $p<.001$ ), with no significant difference in prevalence according to sex (Table 1). Females had more objective snacking occasions than males (2.2 versus. 2.0;  $p<.001$ ) and the most popular snack time period was morning (68.4%), compared to afternoon (59.4%) and late night (31.6%; Table 1). In a model for number of objective snacking occasions with age group, sex, their interaction, energy intake, adiposity (waist circumference or BMI), only sex and energy intake were significant, and the *R*-squared value was only 5%. (Table 1).

Objective snacking contributed 1.8MJ (SD 2.1), or 20.0% (SD 20.0%) of total daily energy intake, and relative to its energy contribution, provided a smaller proportion of daily protein [16.2% (SD 20.6%)], and greater carbohydrate [22.5% (SD 21.0)], total [27.8% (SD 24.9%)] and added sugars [28.3% (SD 31.9%)] (Table 1). The highest proportion of objective snacking energy intake came from the morning snack period (43.8%; Figure 2). Discretionary energy at all three snacking time periods combined contributed to 9.2% of total daily energy intake and 41.3% (SD 37.1%) of snacking energy intake came from discretionary foods (Table 1). Among objective snacking time periods, discretionary foods from the afternoon contributed the most to daily energy

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.



(3.7%, 95% CI 3.3-4.1%) and was the most discretionary-dense (50% of energy intake from discretionary foods) (Figure 3).

Of the top ten food groups contributing to energy intake during snacking, five were core foods: dairy milk, pome fruit, regular breads and bread rolls, nuts and nut products; and tropical and subtropical fruit; and four were discretionary foods: sweet biscuits; cakes, muffins, scones, cake-type desserts; sugar, honey and syrups; and chocolate and chocolate-based confectionery. Coffee and coffee substitutes (the 10<sup>th</sup> food group), which may include both core and discretionary foods, had the largest per-consumer daily grams consumed among the top ten food groups using the objective snacking definition (median 250g, interquartile range: 200–330 g). Dairy milk was the food group that contributed the most to snacking energy intake (mean 7.6%, SD 0.2%) and had the highest prevalence of consumers (30.4%; Table 2).

In a model for the proportion of daily energy from discretionary foods, age group, the interaction of sex and age group, energy intake, and BMI group were all significant and the R-squared was 5.1% (Table 1). Adults classified as having the recommended BMI (18-25kg/m<sup>2</sup>) had a significantly lower proportion of their energy from discretionary foods (41.6%) than those classified as obese (45.4%) ( $p=0.008$ ) but not overweight (41.7%) (Table 1).

Prevalence of snacking was greater using the subjective definition (98.5%) compared to using the objective definition (88.2%), as were the mean frequency of snacking occasions (3.9 versus 2.1 occasions respectively; Table 2). Mean energy intake from subjective snacking was 2.3 MJ (SD 2.2 MJ) which represented 25.8% (SD 18.6%) of

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

total daily energy, compared to a mean of 1.8 MJ (SD 2.1 MJ) and 20.0% (SD 20.0%) of daily energy during objective snacking periods. Using the subjective compared to the objective definition, snacking contributed similar amounts to daily protein, but more to total fat, carbohydrate, total and added sugars (Table 2).

Using the subjective definition, almost half (46.2%) of the discretionary energy came from snacking, compared to 27.0% when objectively defined. Discretionary foods contributed 52.6% (SD 35.2%) of all subjective snacking energy and 41.3% (SD 37.1%) of objective snacking energy. Most of the top food groups were similar across the two snacking definitions, but only in the subjective definition were beers and wines included (Table 2). In models measuring the proportion of snacking energy from discretionary foods adjusted for age group, sex, their interaction, energy intake, and BMI group; for both snacking definitions age group and energy intake were significant, but not BMI group, and the R-squared was 2% for the objective definition and 3% for the subjective definition.

## Discussion

Snacking behaviour was highly prevalent among a representative sample of Australian adults, particularly during the morning. The prevalence of snacking and the contribution to discretionary energy was greater when the subjective definition rather than objective definition was applied. This may be because people perceive discretionary foods as snacks, rather than actual between meal snacking. When objectively defined, snacking contributed more to core food group energy than discretionary energy and the discretionary energy from snacking was not associated with BMI, while overall discretionary energy intake was.

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

Australian snacking prevalence and its contribution to total energy intake is generally consistent with other developed countries<sup>5,8,29,30</sup>; however, snacking behaviour patterns differ, possibly due to cultural differences. In Australia, snacking was most prevalent in the morning; whereas in the USA, Mexico, Brazil, and China, afternoon and late night snacking were the most prevalent<sup>1-3,30,31</sup>. This finding demonstrates the need for nation-specific public health messages and guidelines on snacking given its high prevalence, but also the potential for contribution to the intake of core food groups. Using the objective definition, this study found that under half of snacking energy came from discretionary foods, but more than half, when the subjective definition was used. This demonstrates that snacking is not synonymous with the intake of discretionary foods but may be perceived to be so. Snacking is a prominent dietary pattern in Australia, that has increased over time in frequency and energy contribution<sup>25</sup>. Since 67-97% of Australian adults fall short of dietary recommendations for each of the five core foods groups<sup>32</sup>, there is potential for snacking to help address this national dietary shortfall.

The finding that dairy, fruit and breads are some of the leading food groups that contributed to snacking is consistent with other data globally<sup>3,14,33</sup>, and suggests that reducing snacking could have potential negative implications for the consumption of these food groups and the nutrients they provide. Given that 44% of objective snacking energy was derived from discretionary foods and that snacking contributed more to daily added sugars intake than to daily energy intake, there is still an opportunity to improve the nutritional quality of snacking. Snacking recommendations may need to

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

focus on replacing discretionary foods with snack foods from the five core food groups, rather than limiting snacking entirely.

Using the subjective definition, this study found snacking frequency was high and contributed 54% to total added sugars. This is consistent with a previous study in Australian men that reported a positive association between snacking frequency and added sugars intake <sup>34</sup>. Interventions that promoted fruit for snacking have demonstrated improvements in the total dietary quality, including a reduction in added sugars <sup>35,36</sup> and reduced unhealthy snacking <sup>37</sup>. This may be particularly relevant for the afternoon snacking period, which was the most discretionary dense snacking time period.

A number of studies have suggested that discretionary foods correlated with total energy intakes, both across the day <sup>38,39</sup> and longitudinally over time <sup>40,41</sup>. This study highlights that whilst snacking is often regarded as a top contributor of discretionary food intakes, the high discretionary energy intake among Australian adults is a concern across both main meals and snacking eating occasions. Interventions aimed at reducing discretionary food intake should consider the intake across the entire day, prioritising the evening meal which had the highest total discretionary energy of any eating time period.

Interestingly, whilst the proportion of energy from total discretionary foods was associated with BMI, the proportion of energy from discretionary foods during snacking and the number of snacking occasions, was not. It is possible that the contribution of discretionary food intake from snacking, at 27% of total discretionary

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

energy intake, is insufficient to be independently associated with BMI. The literature on the association between eating occasions and anthropometric measures is consistent with the current findings, with literature reviews generally reporting that eating and snacking frequency are not uniquely associated with body weight <sup>15,42-44</sup>. There may also be compensation for energy consumed in snacks by consuming less energy at main meals, as seen in recent research <sup>45</sup>. A cluster analysis of the snacking patterns of American adults found that different patterns of snacking had different associations with both BMI and waist circumference <sup>46</sup>. Associations between discretionary energy, snacking and adiposity measures in cross-sectional studies may also be explained by confounding or reverse causality. A universal definition for snacking and subsequent clinical trials is required to better understand the relationship between snacking and anthropometry, with consideration to these complexities, before authoritative recommendations on snacking and weight management can be established.

Interestingly, beer and wines were only among the leading food groups when the subjective definition was applied. This may be due to these discretionary foods being consumed during main mealtimes but subjectively regarded as a snack. It is also likely that these foods and beverages are the main reason why discretionary energy differed between subjective and objective snacking, since the other leading food groups for objective and subjective snacking were similar. Core foods eaten during snack periods, when defined objectively, may have also been regarded as a main meal not as snacking, such as dairy milk or bread. Regular breads and bread rolls were the fifth leading food group contributor to total energy from snacking when defined

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

objectively, but outside the top ten when defined subjectively. It has been reported that healthy foods were more likely to be viewed as meals than as snack foods <sup>47</sup>. Given the cognitive bias for snacking with discretionary energy and subsequent food intake, associations between self-reported snacking and diet quality should be interpreted with caution. The findings also suggest that eating occasion-based recommendations may need to clearly define snacking or be in the context of the time of day.

Limitations of the current study include that the objective method does not account for diets that do not fit this assumed pattern of eating <sup>48</sup>, such as those of shift workers or with different cultural eating practices. Cross-sectional data are observational, precluding causal conclusions to be drawn. By nature, dietary recall is often affected by under-reporting, which can influence the associations for eating patterns <sup>49</sup>; however, under-reporters were not excluded as doing so may lead to selection bias and inflated associations, particularly when investigating dietary patterns <sup>50,51</sup>. The reported eating occasion 'beverage/drink' was included in the subjective definition. Future research could include 'beverage/drink' as a separate category from both snacking and main meals to determine if the current findings are specific to foods or beverages. Finally, there are opportunities for future research to perform multivariate analyses to explore what other factors (e.g. behavioural, health, environmental and socioeconomic), and how the interrelationship of these factors, affect snacking habits and discriminate between objective and subjective snacking.

Snacking eating behaviour was highly prevalent among Australian adults, comprised of both core and discretionary foods, and contributed more than a 20% of total energy

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

intake. The contribution of snacking to discretionary intake was associated with its definition, where subjectively defined it contributed more than half of total discretionary intake, suggesting that adults are more likely to perceive discretionary foods as snacks. Associations between self-reported snacking should be interpreted with caution as self-reported snacking is associated with a cognitive bias towards discretionary energy that may not reflect the time of consumption.

**Conflict of interest:** FFM, AM, TC, KT, MB, MW, PP, and SM independently work for Nutrition Research Australia, which is gains funding for projects from government, not-for-profits, professional, community, and industry organisations. FFM, AM, TC, KT, MB, MW, PP and SM declare no conflicts of interest. The funding body, Nestlé Australia Ltd had no contribution to the draft analysis plan, the data analysis, or drafting of the manuscript.

**Funding:** This project has been funded by a research grant from Nestlé Australia Ltd.

**Authorship:** All authors have reviewed and approved the final version of this manuscript prior to submission and declare this work has not be submitted for publication elsewhere. FM led the drafting of the manuscript. AM and PP conducted the data analysis. All authors contributed to the revision of the manuscript and study concept

**Acknowledgements:** The authors would like to acknowledge Dr Jean Kim, Nestlé Australia, who provided top line suggestions on the proposed manuscript.

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

## References

1. Wang Z, Zhai F, Zhang B, Popkin BM. Trends in Chinese snacking behaviors and patterns and the social-demographic role between 1991 and 2009. *Asia Pac J Clin Nutr.* 2012;21(2):253-262.
2. Duffey KJ, Pereira RA, Popkin BM. Prevalence and energy intake from snacking in Brazil: analysis of the first nationwide individual survey. *European journal of clinical nutrition.* 2013;67(8):868-874.
3. Duffey KJ, Rivera JA, Popkin BM. Snacking is prevalent in Mexico. *J Nutr.* 2014;144(11):1843-1849.
4. Vatanparast H, Islam N, Patil RP, Shafiee M, Smith J, Whiting S. Snack Consumption Patterns among Canadians. *Nutrients.* 2019;11(5).
5. Piernas C, Popkin BM. Snacking increased among U.S. adults between 1977 and 2006. *The Journal of nutrition.* 2010;140(2):325-332.
6. Potter M, Vlassopoulos A, Lehmann U. Snacking Recommendations Worldwide: A Scoping Review. *Adv Nutr.* 2018;9(2):86-98.
7. Ovaskainen ML, Reinivuo H, Tapanainen H, Hannila ML, Korhonen T, Pakkala H. Snacks as an element of energy intake and food consumption. *European journal of clinical nutrition.* 2006;60(4):494-501.
8. Dunford EK, Popkin BM. Disparities in Snacking Trends in US Adults over a 35 Year Period from 1977 to 2012. *Nutrients.* 2017;9(8).
9. Murakami K, Livingstone MB. Associations between meal and snack frequency and diet quality and adiposity measures in British adults: findings from the National Diet and Nutrition Survey. *Public Health Nutr.* 2015:1-11.
10. Marmonier C, Chapelot D, Fantino M, Louis-Sylvestre J. Snacks consumed in a nonhungry state have poor satiating efficiency: influence of snack composition on substrate utilization and hunger. *Am J Clin Nutr.* 2002;76(3):518-528.
11. Hibi M, Masumoto A, Naito Y, et al. Nighttime snacking reduces whole body fat oxidation and increases LDL cholesterol in healthy young women. *Am J Physiol Regul Integr Comp Physiol.* 2013;304(2):R94-R101.
12. Mattes RD. Snacking: A cause for concern. *Physiology & Behavior.* 2018;193:279-283.
13. Viskaal-van Dongen M, Kok FJ, de Graaf C. Effects of snack consumption for 8 weeks on energy intake and body weight. *Int J Obes (Lond).* 2010;34(2):319-326.

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.



14. Zizza CA, Xu B. Snacking is associated with overall diet quality among adults. *Journal of the Academy of Nutrition and Dietetics*. 2012;112(2):291-296.
15. Johnson GH, Anderson GH. Snacking definitions: impact on interpretation of the literature and dietary recommendations. *Crit Rev Food Sci Nutr*. 2010;50(9):848-871.
16. Hess JM, Slavin JL. The benefits of defining “snacks”. *Physiology & Behavior*. 2018;193:284-287.
17. Australian Bureau of Statistics. National Health Survey: First Results 2014-15. Cat. No. 4364.0.55.001. In. Canberra: ABS; 2015.
18. Australian Bureau of Statistics. Australian Health Survey: Nutrition First Results - Food and Nutrients, 2011-12; ABS cat no. 4364.0.55.007. In. Canberra: ABS; 2014.
19. Fayet-Moore F, McConnell A, Cassettari T, Tuck K, Petocz P, Kim J. Discretionary intake among Australian adults: prevalence of intake, top food groups, time of consumption and its association with sociodemographic, lifestyle and adiposity measures. *Public Health Nutr*. 2019;22(9):1576-1589.
20. Von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Prev Med*. 2007;45(4):247-251.
21. Australian Health Survey: Users'Guide, 2011-13. In: Statistics ABo, ed. Canberra: Australian Bureau of Statistics; 2013.
22. Hart CN, Raynor HA, Osterholt KM, Jelalian E, Wing RR. Eating and activity habits of overweight children on weekdays and weekends. *Int J Pediatr Obes*. 2011;6(5-6):467-472.
23. Yates L, Warde A. The evolving content of meals in Great Britain. Results of a survey in 2012 in comparison with the 1950s. *Appetite*. 2015;84:299-308.
24. An R. Weekend-weekday differences in diet among U.S. adults, 2003-2012. *Annals of epidemiology*. 2016;26(1):57-65.
25. Fayet-Moore F, Peters V, McConnell A, Petocz P, Eldridge AL. Weekday snacking prevalence, frequency, and energy contribution have increased while foods consumed during snacking have shifted among Australian children and adolescents: 1995, 2007 and 2011-12 National Nutrition Surveys. *Nutr J*. 2017;16(1):65.
26. Food Standards Australia New Zealand. AUSNUT 2011-13- Australian Food, Supplement and Nutrient Database for Estimation of Population Nutrient Intakes. In. Canberra: FSANZ; 2014.

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

27. National Health and Medical Research Council. Australian Dietary Guidelines. In. Canberra: NHMRC; 2013.
28. World Health Organisation. Obesity: Preventing and Managing the Global Epidemic. (Technical Report Series 894). Report of a WHO Consultation. 2000; [http://www.who.int/nutrition/publications/obesity/WHO\\_TRS\\_894/en/](http://www.who.int/nutrition/publications/obesity/WHO_TRS_894/en/). Accessed 4th February 2015.
29. Garriguet D. Canadians' eating habits. *Health Rep.* 2007;18(2):17-32.
30. Kant AK. Eating patterns of US adults: Meals, snacks, and time of eating. *Physiol Behav.* 2018;193(Pt B):270-278.
31. Kant AK, Graubard BI. 40-year trends in meal and snack eating behaviors of American adults. *J Acad Nutr Diet.* 2015;115(1):50-63.
32. Australian Bureau of Statistics (ABS). Australian Health Survey: Consumption of Food Groups from the Australian Dietary Guidelines, 2011-12. ABS Cat. No. 4364.0.55.012. . In. Canberra 2016.
33. Hampl JS, Heaton CL, Taylor CA. Snacking patterns influence energy and nutrient intakes but not body mass index. *Journal of human nutrition and dietetics : the official journal of the British Dietetic Association.* 2003;16(1):3-11.
34. Leech RM, Livingstone KM, Worsley A, Timperio A, McNaughton SA. Meal Frequency but Not Snack Frequency Is Associated with Micronutrient Intakes and Overall Diet Quality in Australian Men and Women. *J Nutr.* 2016;146(10):2027-2034.
35. Beets MW, Turner-McGrievy B, Weaver RG, et al. Intervention leads to improvements in the nutrient profile of snacks served in afterschool programs: a group randomized controlled trial. *Transl Behav Med.* 2016;6(3):329-338.
36. Alinia S, Lassen AD, Krogholm KS, Christensen T, Hels OH, Tetens I. A workplace feasibility study of the effect of a minimal fruit intervention on fruit intake. *Public Health Nutr.* 2011;14(8):1382-1387.
37. Tak NI, Te Velde SJ, Singh AS, Brug J. The effects of a fruit and vegetable promotion intervention on unhealthy snacks during mid-morning school breaks: results of the Dutch Schoolgruitem Project. *J Hum Nutr Diet.* 2010;23(6):609-615.
38. Kant AK. Consumption of energy-dense, nutrient-poor foods by adult Americans: nutritional and health implications. The third National Health and Nutrition Examination Survey, 1988-1994. *Am J Clin Nutr.* 2000;72(4):929-936.

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

39. Rangan AM, Schindeler S, Hector DJ, Gill TP, Webb KL. Consumption of 'extra' foods by Australian adults: types, quantities and contribution to energy and nutrient intakes. *European journal of clinical nutrition*. 2009;63(7):865-871.
40. Popkin BM, Adair LS, Ng SW. Global nutrition transition and the pandemic of obesity in developing countries. *Nutr Rev*. 2012;70(1):3-21.
41. Popkin BM, Gordon-Larsen P. The nutrition transition: worldwide obesity dynamics and their determinants. *Int J Obes Relat Metab Disord*. 2004;28 Suppl 3:S2-9.
42. Raynor HA, Goff MR, Poole SA, Chen G. Eating Frequency, Food Intake, and Weight: A Systematic Review of Human and Animal Experimental Studies. *Front Nutr*. 2015;2:38.
43. Palmer MA, Capra S, Baines SK. Association between eating frequency, weight, and health. *Nutr Rev*. 2009;67(7):379-390.
44. Palmer MA, Capra S, Baines SK. To snack or not to snack: what should we advise for weight management? *Nutr Diet*. 2011;68(1):60-64.
45. Kant AK, Graubard BI. Within-person compensation for snack energy by US adults, NHANES 2007-2014. *Am J Clin Nutr*. 2019;109(4):1145-1153.
46. Nicklas TA, O'Neil CE, Fulgoni VL, 3rd. Snacking patterns, diet quality, and cardiovascular risk factors in adults. *BMC Public Health*. 2014;14:388.
47. Wansink B, Payne CR, Shimizu M. "Is this a meal or snack?" Situational cues that drive perceptions. *Appetite*. 2010;54(1):214-216.
48. Leech RM, Worsley A, Timperio A, McNaughton SA. Temporal eating patterns: a latent class analysis approach. *Int J Behav Nutr Phys Act*. 2017b;14(1):3.
49. Leech RM, Worsley A, Timperio A, McNaughton SA. The role of energy intake and energy misreporting in the associations between eating patterns and adiposity. *European journal of clinical nutrition*. 2017a.
50. Jessri M, Lou WY, L'Abbé MR. Evaluation of different methods to handle misreporting in obesity research: evidence from the Canadian national nutrition survey. *Br J Nutr*. 2016;115(1):147-159.
51. Vainik U, Konstabel K, Lätt E, Mäestu J, Purge P, Jürimäe J. Diet misreporting can be corrected: confirmation of the association between energy intake and fat-free mass in adolescents. *Br J Nutr*. 2016;116(8):1425-1436.
52. *Australian Dietary Guidelines*. Canberra: NHMRC, Australian Government;2013.

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

## Figure legends

**FIGURE 1:** Percent of energy consumed by time of day and classification of meal and snack time periods as follows: breakfast (5.30am – 9.30am), morning snack (9.30am – 11.30am), midday meal (11.30am – 2.30pm), afternoon snack (2.30pm – 5.30pm), evening meal (5.30pm – 9.30pm), late night snack (9.30pm – 5.30am). Data from adults 19+y from the 2011-12 National Nutrition and Physical Activity Survey.

**FIGURE 2:** Based on the objective definition of snacking: per capita contribution of main meals and snacking periods to total energy intake (percent of total energy with 95% confidence intervals shown, and mean energy intake) among Australian adults 19+y. Main meals occurred between 5.30am – 9.30am, 11.30am – 2.30pm, and 5.30pm – 9.30pm. Snacking periods occurred between 9.30am – 11.30am (morning snack), 2.30pm – 5.30pm (afternoon snack), and 9.30pm – 5.30am (late night snack). Data from the 2011-12 National Nutrition and Physical Activity Survey.

**FIGURE 3:** Based on the objective definition of snacking: per capita contribution of main meals and snacking periods to total energy intake by core versus discretionary foods among Australian adults 19+y (error bars show 95% confidence intervals). Main meals and snacking periods defined as breakfast (5.30am – 9.30am), morning snack (9.30am – 11.30am), midday meal (11.30am – 2.30pm), afternoon snack (2.30pm – 5.30pm), evening meal (5.30pm – 9.30pm), late night snack (9.30pm – 5.30am). Data from the 2011-12 National Nutrition and Physical Activity Survey.

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

**Table 1:** Descriptive characteristics and snacking <sup>(a)</sup> patterns of 8,361 adults 19+y from the 2011-12 National Nutrition and Physical Activity Survey

| Characteristic   | Results            | P-value |
|--|--------------------|---------|
| <b>Females (%)</b>   | 51.1               |         |
| <b>Age group (%)</b>   |                    |         |
| 19-30y   | 22.8               |         |
| 31-50y   | 37.0               |         |
| 51-70y   | 29.2               |         |
| 71+y   | 11.0               |         |
| <b>Weight status (%) <sup>(b)</sup></b>  |                    |         |
| Underweight  | 1.8                |         |
| Normal weight  | 35.7               |         |
| Overweight   | 36.4               |         |
| Obese  | 26.1               |         |
| <b>Waist circumference category (%) <sup>(c)</sup></b>                               |                    |         |
| Not at risk of metabolic complications   | 37.5               |         |
| Increased risk of metabolic complications  | 23.0               |         |
| Greatly increased risk of metabolic complications                                    | 39.5               |         |
| <b>Prevalence of snacking (%) <sup>(d)</sup></b>                                     | 88.2 [87.5 – 88.9] |         |
| <b>Prevalence of snacking by sex (%) <sup>(d)</sup></b>                              |                    | 0.11    |
| Males (n=4092)   | 87.7 [86.7 – 88.7] |         |
| Females (n=4269)   | 88.8 [87.8 – 89.7] |         |
| <b>Prevalence of snacking by age group (%) <sup>(d)</sup></b>                        |                    | <0.001  |
| 19-30y (n=1909)  | 87.7 [86.2 – 89.1] |         |
| 31-50y (n=3091)  | 89.8 [88.8 – 90.9] |         |
| 51-70y (n=2440)  | 88.0 [86.7 – 89.3] |         |
| 71+y (n=921)   | 84.8 [82.5 – 87.1] |         |
| <b>Prevalence of consumption at each meal and snacking period (%) <sup>(d)</sup></b> |                    | <0.001  |
| Breakfast  | 89.4 [88.7 - 90.1] |         |
| Morning snack  | 68.4 [67.4 - 69.4] |         |
| Midday meal  | 87.4 [86.6 - 88.1] |         |
| Afternoon snack  | 59.4 [58.4 - 60.5] |         |
| Evening meal   | 97.0 [96.6 - 97.4] |         |

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. Nutrition and Dietetics, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

|  |                                    |        |
|--|------------------------------------|--------|
| Late night snack   | 31.6 [30.6 - 32.6]                 |        |
| <b>Frequency of snacking (%)</b>   |                                    |        |
| Non-snack consumers (0 snacking occasions)   | 11.8                               |        |
| 1 snacking occasion  | 25.2                               |        |
| 2 snacking occasions   | 29.8                               |        |
| 3 snacking occasions   | 18.4                               |        |
| 4+ snacking occasions  | 14.8                               |        |
| <b>Number of snacking occasions <sup>(e)</sup></b>   | 2.1 (1.4)                          |        |
| <b>Number of snacking occasions by sex <sup>(e)</sup></b>  |                                    | 0.093  |
| Males  | 2.1 (1.4)                          |        |
| Females  | 2.1 (1.4)                          |        |
| <b>Number of snacking occasions by sex <sup>(f, g)</sup></b>                                     |                                    | <0.001 |
| Males  | 2.0 [1.9 – 2.1]                    |        |
| Females  | 2.2 [2.1 – 2.3]                    |        |
| <b>Number of snacking occasions by age group (mean) <sup>(e)</sup></b>                           |                                    | 0.015  |
| 19-30y   | 2.1 (1.5)                          |        |
| 31-50y   | 2.1 (1.4)                          |        |
| 51-70y   | 2.1 (1.4)                          |        |
| 71+y   | 2.0 (1.4)                          |        |
| <b>Proportion of total energy from discretionary foods by BMI group (%) <sup>(f, h)</sup></b>    |                                    | <0.001 |
| Underweight  | 32.5 [29.0 – 36.0] <sup>*,**</sup> |        |
| Normal weight  | 29.8 [29.0 – 30.6] <sup>*</sup>    |        |
| Overweight   | 32.6 [31.8 – 33.4] <sup>*</sup>    |        |
| Obese  | 33.5 [32.6 – 34.5] <sup>*</sup>    |        |
| <b>Proportion of snacking energy from discretionary foods by BMI group (%) <sup>(f, i)</sup></b> |                                    | 0.008  |
| Underweight  | 44.7 [37.6 – 51.8]                 |        |
| Normal weight  | 41.6 [39.9 – 43.2]                 |        |
| Overweight   | 41.7 [40.0 – 43.3]                 |        |
| Obese  | 45.4 [43.4 – 47.3]                 |        |
| <b>Total energy intake (MJ) <sup>(e)</sup></b>   | 8.5 (3.6)                          |        |
| <b>Energy during snacking (MJ) <sup>(e)</sup></b>  | 1.8 (2.1)                          |        |
| <b>Energy contribution of snacking to total energy intake <sup>(e)</sup></b>                     | 20.0 (20.0)                        |        |

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. Nutrition and Dietetics, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

**Contribution from snacking to total macronutrient intake <sup>(e)</sup>**

|  |                    |
|--|--------------------|
| Protein  | 16.2 (20.6)        |
| Fat  | 19.8 (22.9)        |
| Carbohydrate   | 22.5 (21.0)        |
| Total sugars   | 27.8 (24.9)        |
| Added sugars   | 28.3 (31.9)        |
| <b>Total discretionary <sup>(i)</sup> energy intake (MJ) <sup>(e)</sup></b>  | <b>2.9 (2.5)</b>   |
| <b>Contribution of snacking to total discretionary energy <sup>(e)</sup></b> | <b>27.0 (31.4)</b> |
| <b>Energy from discretionary food during snacking <sup>(e)</sup></b>         | <b>41.3 (37.1)</b> |

- 
- a. Based on the objective definition of snacking which uses time of day to define main meals and snacking periods. Snacking periods occurred between 9.30am – 11.30am (morning snack), 2.30pm – 5.30pm (afternoon snack), and 9.30pm – 5.30am (late night snack)
  - b. Based on BMI: underweight (<18.5), normal weight (≥18.5, <25.0), overweight (≥25.0, <30.0), obese (≥30.0).
  - c. Based on World Health Organization cut-offs for waist circumference: not at risk of metabolic complications (females: <80cm; males: <94cm); increased risk of metabolic complications (females: ≥80cm, <88cm; males: ≥94cm, <102cm); substantially increased risk of metabolic complications (females: ≥88cm; males: ≥102cm) <sup>28</sup>.
  - d. Data are expressed as % [95% confidence interval].
  - e. Data are expressed as mean (standard deviation).
  - f. Data are expressed as mean [95% confidence interval].
  - g. Adjusted for age group, their interaction, waist circumference, BMI group, and energy intake.
  - h. Adjusted for sex, age group, their interaction, BMI group, and energy intake. Age group, the interaction of sex and age group, energy intake, and BMI group were significant in the model, R-squared = 0.051; p<0.001 for all.
  - i. Adjusted for sex, age group, their interaction, BMI group and energy intake. Age group and energy intake were significant in the model, R-squared = 0.019; p<0.001 for all.
  - j. Discretionary foods are defined by the Australian Dietary Guidelines as foods and beverages not necessary for a healthy diet and are generally high in saturated fat and/or added sugars, added salt or alcohol and low in fibre <sup>52</sup>.

<sup>\*,\*\*</sup> Different superscript Asterix denote significant difference between groups (p<0.001) by post hoc, Bonferroni.

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

**Table 2:** Snacking patterns and comparison of the top 10 sub-major food and beverage groups by percent energy contribution to snacking among 8,361 Australian adults 19y+ from the 2011-12 National Nutrition and Physical Activity Survey based on the objective and subjective snacking definitions

| Characteristic   | Objective definition <sup>(a)</sup> | Subjective definition <sup>(b)</sup> |
|--|-------------------------------------|--------------------------------------|
| <b>Prevalence of snacking (%) <sup>(c)</sup></b>                                   | 88.2 [87.5 – 88.9]                  | 98.5 [98.2 – 98.8] *                 |
| <b>Frequency of snacking (%)</b>   |                                     |                                      |
| Non-snack consumer (0 snacking occasions)  | 11.8                                | 1.5                                  |
| 1 snacking occasion  | 25.2                                | 8.2                                  |
| 2 snacking occasions   | 29.8                                | 15.4                                 |
| 3 snacking occasions   | 18.4                                | 21.4                                 |
| 4+ snacking occasions  | 14.8                                | 53.5                                 |
| <b>Number of snacking occasions <sup>(d)</sup></b>                                 | 2.1 (1.4)                           | 3.9 (2.1)                            |
| <b>Energy during snacking (MJ) <sup>(d)</sup></b>                                  | 1.8 (2.1)                           | 2.3 (2.2) *                          |
| <b>Energy contribution of snacking to total energy intake (%) <sup>(d)</sup></b>   | 20.0 (20.0)                         | 25.8 (18.6) *                        |
| <b>Contribution from snacking to total macronutrient intake (%) <sup>(d)</sup></b> |                                     |                                      |
| Protein  | 16.2 (20.6)                         | 16.0 (16.3) **                       |
| Fat  | 19.8 (22.9)                         | 23.2 (21.4) *                        |
| Carbohydrate   | 22.5 (21.0)                         | 29.5 (20.6) *                        |
| Total sugars   | 27.8 (24.9)                         | 42.7 (27.0) *                        |
| Added sugars   | 28.3 (31.9)                         | 46.8 (35.9) *                        |
| <b>Contribution of snacking to total discretionary energy (%) <sup>(d)</sup></b>   | 27.0 (31.4)                         | 46.2 (34.7) *                        |
| <b>Energy from discretionary food during snacking (%) <sup>(d)</sup></b>           | 41.3 (37.1)                         | 52.6 (35.2) *                        |
| <b>Total energy from snacking (%) <sup>(e)</sup></b>                               |                                     |                                      |

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.



|  |                    |                    |
|--|--------------------|--------------------|
| Dairy milk (cow, sheep, goat)                            | 7.6 ± 0.2          | 8.1 ± 0.2          |
| Sweet biscuits   | 6.5 ± 0.2          | 6.6 ± 0.2          |
| Coffee and coffee substitutes                            | 5.8 ± 0.2          | 5.5 ± 0.2          |
| Cakes, muffins, scones, cake-type desserts               | 5.7 ± 0.2          | 5.9 ± 0.2          |
| Pome fruit   | 5.7 ± 0.2          | 4.7 ± 0.2          |
| Regular breads and bread rolls (plain/unfilled/untopped) | 5.0 ± 0.2          | – <sup>g</sup>     |
| Sugar, honey, and syrups                                 | 3.6 ± 0.1          | 4.0 ± 0.1          |
| Nuts and nut products                                    | 3.3 ± 0.2          | 4.3 ± 0.2          |
| Tropical and subtropical fruit                           | 3.2 ± 0.2          | – <sup>g</sup>     |
| Chocolate and chocolate-based confectionary              | 2.8 ± 0.2          | 4.8 ± 0.2          |
| Beers  | – <sup>g</sup>     | 4.5 ± 0.2          |
| Wines  | – <sup>g</sup>     | 3.0 ± 0.1          |
| <b>Consumers snacking (%) <sup>(f)</sup></b>             |                    |                    |
| Dairy milk (cow, sheep, goat)                            | 30.4 [29.4 - 31.4] | 43.3 [42.3 - 44.4] |
| Sweet biscuits   | 12.8 [12.0 - 13.5] | 19.3 [18.5 - 20.2] |
| Coffee and coffee substitutes                            | 28.9 [27.9 - 29.8] | 43.5 [42.4 - 44.6] |
| Cakes, muffins, scones, cake-type desserts               | 8.3 [7.7 - 8.9]    | 11.1 [10.4 - 11.8] |
| Pome fruit   | 11.3 [10.6 - 12.0] | 15.9 [15.1 - 16.6] |
| Regular breads and bread rolls (plain/unfilled/untopped) | 13.1 [12.4 - 13.8] | – <sup>g</sup>     |
| Sugar, honey, and syrups                                 | 20.6 [19.7 - 21.4] | 29.1 [28.1 - 30.1] |
| Nuts and nut products                                    | 6.7 [6.1 - 7.2]    | 11.5 [10.9 - 12.2] |
| Tropical and subtropical fruit                           | 7.8 [7.2 - 8.4]    | – <sup>g</sup>     |

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

|  |                 |                    |
|--|-----------------|--------------------|
| Chocolate and chocolate-based confectionary              | 6.0 [5.5 - 6.5] | 13.9 [13.1 - 14.6] |
| Beers  | –g              | 9.3 [8.7 - 10.0]   |
| Wines  | –g              | 6.5 [5.9 - 7.0]    |
| <b>Snacking intake (g) <sup>(h)</sup></b>                |                 |                    |
| Dairy milk (cow, sheep, goat)                            | 62 [31 - 146]   | 62 [31 - 146]      |
| Sweet biscuits   | 25 [17 - 37]    | 25 [17 - 38]       |
| Coffee and coffee substitutes                            | 250 [200 - 330] | 281 [200 - 450]    |
| Cakes, muffins, scones, cake-type desserts               | 98 [57 - 163]   | 105 [62 - 163]     |
| Pome fruit   | 164 [164 - 188] | 164 [158 - 188]    |
| Regular breads and bread rolls (plain/unfilled/untopped) | 64 [54 - 80]    | –g                 |
| Sugar, honey, and syrups                                 | 7 [4 - 13]      | 8 [4 - 13]         |
| Nuts and nut products                                    | 28 [14 - 36]    | 28 [15 - 36]       |
| Tropical and subtropical fruit                           | 98 [98 - 111]   | –g                 |
| Chocolate and chocolate-based confectionary              | 33 [15 - 53]    | 30 [15 - 51]       |
| Beers  | –g              | 758 [379 - 1515]   |
| Wines  | –g              | 312 [208 – 594]    |

\* p<0.001; \*\* p=0.497.

- The objective definition of snacking uses time of day to define main meals and snacking periods. Snacking periods occurred between 9.30am – 11.30am (morning snack), 2.30pm – 5.30pm (afternoon snack), and 9.30pm – 5.30am (late night snack).
- The subjective definition of snacking used the self-reported eating occasion provided by respondents during dietary recall. Snacking was defined as any of the following self-reported eating occasions: morning tea, snack, afternoon tea, beverage/drink, supper, other, extended consumption.
- Data are expressed as % [95% confidence interval].
- Data are expressed as mean (standard deviation).
- Data are expressed as mean  $\pm$  standard error.
- Data are expressed as % [95% confidence interval].

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

- 
- g. Was not in the top 10 sub-major food and beverage groups by percent energy contribution to snacking.
  - h. Data are expressed as median [interquartile range].
- 

---

This is the peer reviewed version of the following article:

Fayet-Moore, F., McConnell, A., Petocz, P., Cassettari, T., Tuck, K., Blumfield, M., Warner, M., & Marshall, S. (2020). Contribution of dietary snacking behaviours to discretionary energy intake and anthropometric measures in Australian adults: A comparison using an objective vs subjective definition for snacking. *Nutrition and Dietetics*, which has been published in final form at <https://doi.org/10.1111/1747-0080.12648>.

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.